

Contents

Revision of the concept of the orthide brachiopod <i>Cyrtinotella</i> in the Middle Ordovician of the East Baltic <i>Linda Hints</i>	63
<i>Cruziana</i> traces from the Late Silurian (Pridoli) carbonate shelf of Saaremaa, Estonia <i>Olev Vinn</i>	71
Postglacial occurrence and decline of <i>Betula nana</i> L. (dwarf birch) in northeastern Poland <i>Danuta Drzymulska</i>	76
Ice-free interval corresponding to Marine Isotope Stages 4 and 3 at the Last Glacial Maximum position at Kileshino, Valdaj Upland, Russia <i>Katrin Lasberg, Volli Kalm and Kersti Kihno</i>	88
Geological settings of the protected Selisoo mire (northeastern Estonia) threatened by oil shale mining <i>Helen Hiiemaa, Mario Mustasaar, Marko Kohv, Tiit Hang, Argo Jõeleht, Katrin Lasberg and Volli Kalm</i>	97
Thunderstorms caused by southern cyclones in Estonia <i>Kaupo Mändla, Sven-Erik Enno and Mait Sepp</i>	108

SHORT COMMUNICATION

First record of the trace fossil <i>Oikobesalon</i> from the Ordovician (Darriwilian) of Baltica <i>Olev Vinn and Ursula Toom</i>	118
--	-----